IMM-NYU 314
NOVEMBER 1963

NEW YORK UNIVERSITY
COURANT INSTITUTE OF
MATHEMATICAL SCIENCES

Asymptotic Efficiency of a Class of c-Sample Tests

MADAN LAL PURI

PREPARED UNDER
CONTRACT NONR-285(38)
WITH THE
OFFICE OF NAVAL RESEARCH

IMM-314

C.1

NEW YORK UNIVERSITY
Courant Institute of Mathematical Sciences

ASYMPTOTIC EFFICIENCY OF A CLASS OF c -SAMPLE TESTS¹

Madan Lal Puri

¹This paper was prepared with the partial support of the Office of Naval Research, Contract Nonr-222-(43), while the author was at the University of California, Berkeley. It was revised at the Courant Institute of Mathematical Sciences, New York University under the sponsorship of the Office of Naval Research, Contract Nonr-285(38). Reproduction in whole or in part is permitted for any purpose of the United States Government.

1. Summary. For testing the equality of c continuous probability distributions on the basis of c independent random samples, the test statistics of the form

$$\mathcal{L} = \sum_{j=1}^c m_j \left[(T_{N,j} - \mu_{N,j}) / A_N \right]^2$$

are considered. Here m_j is the size of the j th sample, $\mu_{N,j}$ and A_N are normalizing constants, and

$$T_{N,j} = \frac{1}{m_j} \sum_{i=1}^N E_{N,i} Z_{N,i}^{(j)}$$

where $Z_{N,i}^{(j)} = 1$, if the i th smallest of $N = \sum_{j=1}^c m_j$ observations is from the j th sample and $Z_{N,i}^{(j)} = 0$ otherwise. Sufficient conditions are given for the joint asymptotic normality of $T_{N,j}$; $j=1, \dots, c$. Under suitable regularity conditions and the assumption that the i th distribution function is $F(x + \theta_i / \sqrt{N})$, the limiting distribution of $\mathcal{L}$ is derived. Finally, the asymptotic relative efficiencies in Pitman's sense of the $\mathcal{L}$ test relative to some of its competitors viz, the Kruskal-Wallis H test (which is a particular case of the $\mathcal{L}$ test) and the classical F test are obtained and shown to be independent of the number c of samples.

2. Introduction. One of the frequently encountered problems in statistics is to decide whether differences in various samples should be regarded as due to differences in the parent populations or due to chance variations which are to be expected among random samples from the same population. A few tests of nonparametric nature have been proposed for this c-sample problem. The Kruskal-Wallis H test [14], Terpestra's c-sample test [26], the Mood and Brown c-sample test [22] and Kiefer's K-sample analogues of the Kolmogorov-Smirnov and Cramér-von Mises tests [12] are a few of them. Tests for two-sample problems have been proposed by Wilcoxon [29], Mann and Whitney [19], Mood and Brown [22], Lehmann [15] and others. Consistency and power properties of some of these tests have been discussed by Lehmann [15], [16], [17], Mood [23], Van Dantzig [5] and others. An exhaustive review of this problem is given in Kruskal and Wallis [14] and Scheffé [25].

The H test of Kruskal and Wallis is a direct generalization of the two-sample Wilcoxon test discussed in detail by Mann and Whitney [19], and its limiting distribution has been derived by Kruskal [13] under the null hypothesis and by Andrews [1] under an alternative hypothesis. These results are generalized by those of the present paper concerning the limiting distribution of the L test.

The problem discussed in this paper originated from the paper of Chernoff and Savage [2] and had its basis in the paper of Hodges and Lehmann [10]. In their paper "The efficiency of some nonparametric competitors of the t-test" [10], Hodges and

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

61

62

63

64

Lehmann discussed the asymptotic efficiency of the Wilcoxon test with respect to all translation alternatives. In the same paper they conjectured that the normal score test which was known to be as efficient as the t-test for normal alternatives [35] is at least as efficient as the t-test for all other alternatives. The validity of this conjecture was established by Chernoff and Savage [2], who developed a new theorem for asymptotic normality of normal score test statistics for the two-sample problem and by a variational argument proved the Hodges-Lehmann conjecture. The work presented here is an attempt toward generalizing these results to the c-sample problem.

Formally, we may state the c-sample problem as follows.

Let $\{X_{ij}, j=1, \dots, m_i; i=1, \dots, c\}$ be a set of independent random variables and let $F^{(i)}(x)$ be the probability distribution of X_{ij} . The set of admissible hypotheses designates that each $F^{(i)}(x)$ belongs to some class of distribution functions Ω . The hypothesis to be tested, say H_0 , specifies that $F^{(i)}$ is an element of Ω , for each i , and that furthermore

$$(2.1) \quad F^{(1)}(x) = \dots = F^{(c)}(x) \quad \text{for all real } x.$$

The class of alternatives to H_0 can be considered to be all sets $(F^{(1)}(x), \dots, F^{(c)}(x))$ which belong to Ω but which violate (2.1). To avoid the problem of ties, it is assumed throughout that the class Ω is the class of continuous distribution functions.

After finding sufficient conditions for the joint asymptotic normality of $T_{N,j}$; $j=1,\dots,c$, we study the limiting distributions of $\mathcal{L}$ under sequences of admissible alternative hypothesis H_n^P which specifies that for each $i=1,2,\dots,c$; $F^{(1)}(x) = F(x+\theta_i/\sqrt{n})$ with $F \in \Omega$ but not specified further, and for some pair (i,j) , $\theta_i \neq \theta_j$ where the θ_i 's are real numbers. Limiting probability distributions of $\mathcal{L}$ will then be found as $n \rightarrow \infty$. The problem will be so formulated that $m_i(n)/n$ tends to some limit s_i , $0 < s_i < \infty$, as n tends to ∞ .

3. The proposed test and its relationship to other tests.

The over-all sample consists of $\sum_{i=1}^c m_i = N$ independent random variables X_{ij} ($i=1,\dots,c$; $j=1,\dots,m_i$), where the first subscript refers to the subsample and the second subscript indexes observations within a subsample. Under the null hypothesis all the X 's have the same continuous but unknown c.d.f. (cumulative distribution function) $F(x)$.

Let $Z_{N,i}^{(j)} = 1$, if the i th smallest observation from the combined sample of size N is from the j th sample and otherwise let $Z_{N,i}^{(j)} = 0$. Denote

$$(3.1) \quad m_j T_{N,j} = \sum_{i=1}^N Z_{N,i}^{(j)} E_{N,i}$$

where $E_{N,i}$ are given numbers. Then we propose to consider the test statistic $\mathcal{L}$ defined as

$$(3.2) \quad \mathcal{L} = \sum_{j=1}^c m_j \left[(T_{N,j} - \mu_{N,j}) / A_N \right]^2$$

where $\mu_{N,j}$ and A_N are normalizing constants for the statistics $T_{N,j}$; $j=1, \dots, c$.

The $\mathcal{L}$ test presented in this paper includes as special cases a number of well-known tests. For example, when $E_{N,i} = i/N$, it becomes the Kruskal-Wallis H test which is a direct generalization of the two-sample Wilcoxon test and is related to Terpestra's K -sample test [26]. When $c = 2$ and $E_{N,i}$ is the expected value of the i th order statistic from the standard normal distribution, then the $\mathcal{L}$ test coincides with the symmetrical two-tail version of the normal score test, also known as the Fisher-Yates-Terry-Hoeffding c_1 test and which is asymptotically equivalent to Van der Waerden's test [30], [31]. For it is then seen that

$$\mathcal{L} = \frac{N}{N-m_1} \left[\sum_{i=1}^{m_2} E(V^{(s_i)} | s_1) \right]^2$$

where $V^{(1)} < \dots < V^{(N)}$ is an ordered sample of size N from a standard normal distribution, and $s_1 < \dots < s_{m_2}$ are the ranks of $X_{21}, \dots, X_{2m_2}$ from the combined sample. See Lehmann [17], pp. 236-237. When $c = 2$, and $E_{N,i} = |1/2 - i/N|$, the $\mathcal{L}$ -test reduces to the Freund-Ansari test [8] for testing the equality of dispersion of two populations.

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

4. Assumptions and notations. Let $X_{i1}, \dots, X_{im_i}$ be the ordered observations of a random sample from a population with continuous c.d.f. (cumulative distribution function) $F^{(i)}(x)$; $i=1, \dots, c$. Let $N = \sum_{i=1}^c m_i$ and $\lambda_i = m_i/N$ and assume that for all N , the inequalities $0 < \lambda_0 \leq \lambda_1, \dots, \lambda_c \leq 1 - \lambda_0 < 1$ hold for some fixed $\lambda_0 \leq 1/c$.

Let

$$S_{m_i}^{(i)}(x) = m_i^{-1} \quad (\text{number of } X_{ij} \leq x, j=1, \dots, m_i)$$

be the sample c.d.f. of the m_i observations in the i th set. We shall omit the subscript m_i whenever this causes no confusion.

Define

$$H_N(x) = \lambda_1 S_{m_1}^{(1)}(x) + \dots + \lambda_c S_{m_c}^{(c)}(x).$$

Thus $H_N(x)$ is the combined sample c.d.f. The combined population c.d.f. is

$$H(x) = \lambda_1 F^{(1)}(x) + \dots + \lambda_c F^{(c)}(x).$$

Even though $H(x)$ depends on N through the λ 's, our notation suppresses this fact for convenience and also because our limit theorems are uniform with respect to $F^{(1)}, \dots, F^{(c)}$ and $\lambda_1, \dots, \lambda_c$.

Let $Z_{N,i}^{(j)} = 1$ if the i th smallest of $N = \sum_{i=1}^c m_i$ observations is from the j th set and otherwise let $Z_{N,i}^{(j)} = 0$.

Denote

$$(4.1) \quad \tau_{N,j} = m_j \cdot T_{N,j} = \sum_{i=1}^N E_{N,i} Z_{N,i}^{(j)}$$

where the $E_{N,i}$ are given numbers. Following Chernoff and Savage [2], we shall use the representation

$$(4.2) \quad T_{N,j} = \int_{-\infty}^{\infty} J_N[H_N(x)] dS_{m_j}^{(j)}(x)$$

where $E_{N,i} = J_N(i/N)$. While J_N need be defined only at $1/N, 2/N, \dots, N/N$, we shall find it convenient to extend its domain of definition to $(0,1]$ by letting J_N be constant on $(i/N, (i+1)/N]$.

Let

$$I_N = [x: 0 < H_N(x) < 1].$$

Then I_N is a random interval, given by

$$I_N = [X^{(1)}, X^{(N)}]$$

where $X^{(1)} < \dots < X^{(N)}$ denote the N observations arranged according to size.

Throughout, K will be used as a generic constant which may depend on J_N but will not depend on $F^{(1)}, \dots, F^{(c)}$, $m_1, \dots, m_c$ and N . The methods used in the proof for the asymptotic normality of the $T_{N,j}$'s are mainly adaptations of the methods of Chernoff and Savage [2].

5. Joint asymptotic normality. Before proving the asymptotic normality of the $T_{N,j}$'s we state a few elementary results.

$$(5.1) \quad H \geq \lambda_1 F^{(1)} \geq \lambda_0 F^{(1)}; \quad i=1, \dots, c.$$

$$(5.2) \quad 1 - F^{(1)} \leq \frac{1-H}{\lambda_1} \leq \frac{1-H}{\lambda_0}; \quad i=1, \dots, c.$$

$$(5.3) \quad F^{(1)}(1-F^{(1)}) \leq \frac{H(1-H)}{\lambda_1^2} \leq \frac{H(1-H)}{\lambda_0^2}; \quad i=1, \dots, c.$$

$$(5.4) \quad dH \geq \lambda_1 dF^{(1)} \geq \lambda_0 dF^{(1)}; \quad i=1, \dots, c.$$

Lemma 5.1. If

(1) $J(H) = \lim_{N \rightarrow \infty} J_N(H)$ exists for $0 < H < 1$ and is not
constant,

$$(2) \quad \int_{I_N} [J_N(H_N) - J(H_N)] dS_{m_j}^{(j)}(x) = o_p(N^{-(1/2)}),$$

$$(3) \quad J_N(1) = o(\sqrt{N})$$

$$(4) \quad \left| J^{(i)}(H(x)) \right| = \left| \frac{d^i J(H)}{dH^i} \right| \leq K[H(1-H)]^{-i-(1/2)+\delta}$$

for $i=0,1,2$, and for some $\delta > 0$,
and almost all x (a.a.x),

then, for fixed $F^{(1)}, \dots, F^{(c)}$ and $\lambda_1, \dots, \lambda_c$,

$$(5.5) \quad \lim_{N \rightarrow \infty} P \left(\frac{T_{N,j} - \mu_{N,j}}{\sigma_{N,j}} \leq t \right) = \int_{-\infty}^t \frac{1}{\sqrt{2\pi}} e^{-x^2/2} dx,$$

where

$$(5.6) \quad \mu_{N,j} = \int_{-\infty}^{+\infty} J[H(x)] dF^{(j)}(x)$$

and

$$(5.7) \quad N \sigma_{N,j}^2 = 2 \sum_{\substack{i=1 \\ i \neq j}}^c \lambda_i \iint_{-\infty < x < y < \infty} F^{(i)}(x) [1 - F^{(i)}(y)] J'[H(x)] J'[H(y)] dF^{(j)}(x) dF^{(j)}(y) \\ + \frac{2}{\lambda_j} \sum_{\substack{i=1 \\ i \neq j}}^c \lambda_i^2 \iint_{-\infty < x < y < \infty} F^{(j)}(x) [1 - F^{(j)}(y)] J'[H(x)] J'[H(y)] dF^{(i)}(x) dF^{(i)}(y) \\ + \frac{1}{\lambda_j} \sum_{\substack{i,k=1 \\ i \neq k \\ i \neq j \\ k \neq j}}^c \lambda_i \lambda_k \left[\iint_{-\infty < x < y < \infty} F^{(j)}(x) [1 - F^{(j)}(y)] J'[H(x)] J'[H(y)] dF^{(i)}(x) dF^{(k)}(y) \right. \\ \left. + \iint_{-\infty < y < x < \infty} F^{(j)}(y) [1 - F^{(j)}(x)] J'[H(x)] J'[H(y)] dF^{(i)}(x) dF^{(k)}(y) \right].$$

Proof.

$$T_{N,j} = \int_{x=-\infty}^{x=+\infty} J_N[H_N(x)] dS_{m_j}^{(j)}(x) \\ = \int_{[x: 0 < H_N(x) < 1]} [J_N[H_N(x)] - J[H_N(x)]] dS_{m_j}^{(j)}(x) \\ + \int_{[x: 0 < H_N(x) < 1]} J[H_N(x)] dS_{m_j}^{(j)}(x) + \int_{[x: H_N(x)=1]} J_N[H_N(x)] dS_{m_j}^{(j)}(x).$$

In the second integral, writing $dS_{m_j}^{(j)}(x) = d(S_{m_j}^{(j)}(x) - F^{(j)}(x) + F^{(j)}(x))$,

$$J[H_N(x)] = J[H(x)] + [H_N(x) - H(x)]J'[H(x)] + \frac{[H_N(x) - H(x)]^2}{2} J''[\theta H_N(x) + (1-\theta)H(x)], \text{ a.a.x.},$$

where $0 < \theta < 1$; and $H(x) = \sum_{i=1}^c \lambda_i F^{(i)}(x)$, and simplifying, we obtain

$$T_{N,j} = A + B_{1N} + B_{2N} + \sum_{i=1}^{c+4} C_{iN}$$

where

$$(5.8) \quad A = \int_{[x: 0 < H(x) < 1]} J[H(x)] dF^{(j)}(x)$$

$$(5.9) \quad B_{1N} = \int_{[x: 0 < H(x) < 1]} J[H(x)] d[S_{m_j}^{(j)}(x) - F^{(j)}(x)]$$

$$(5.10) \quad B_{2N} = \int_{[x: 0 < H(x) < 1]} [H_N(x) - H(x)] J'[H(x)] dF^{(j)}(x)$$

(5.11)

$$C_{1,N} = \lambda_1 \int_{[x: 0 < H(x) < 1]} \left[S_{m_1}^{(1)}(x) - F^{(1)}(x) \right] J'[H(x)] d \left[S_{m_j}^{(j)}(x) - F^{(j)}(x) \right]$$

$i=1, \dots, c.$

$$(5.12) \quad C_{c+1,N} = \int_{I_N} \frac{[H_N(x) - H(x)]^2}{2} J''[\theta H_N + (1-\theta)H] dS_{m_j}^{(j)}(x).$$

$$(5.13) \quad C_{c+2,N} = \int_{I_N} \left[J_N[H_N(x)] - J[H_N(x)] \right] dS_{m_j}^{(j)}(x).$$

$$(5.14) \quad C_{c+3,N} = \int_{H_N=1} J_N[H_N(x)] dS_{m_j}^{(j)}(x).$$

$$(5.15) \quad C_{c+4,N} = \int_{H_N=1} \left[-J[H(x)] - [H_N(x) - H(x)] J'[H(x)] \right] dS_{m_j}^{(j)}(x).$$

The proof of the lemma is accomplished by showing that (i) the A-term is nonrandom and finite, (ii) $B_{1N} + B_{2N}$ has a Gaussian distribution in the limit and (iii) the C terms are of higher order.

That the term

$$A = \int_{[x: 0 < H(x) < 1]} J[H(x)] dF^{(j)}(x)$$

is finite and nonrandom follows from assumption 4 of lemma 5.1; see also in this connection [2], p. 986, and in the appendix we have shown that the C terms are of higher order. Thus, all that is required is to prove.

Sub-lemma 5.1. $B_{1N} + B_{2N}$ has a Gaussian distribution in the limit.

Proof. Integrating B_{2N} by parts, replacing $H_N(x) - H(x)$ by $\sum_{i=1}^c \lambda_i [S_{m_i}^{(i)}(x) - F^{(i)}(x)]$, and adding B_{1N} to it, we obtain

$$(5.16) \quad B_{1N} + B_{2N} = - \sum_{\substack{i=1 \\ i \neq j}}^c \lambda_i \int_{x=-\infty}^{x=+\infty} B(x) d \left[S_{m_i}^{(i)}(x) - F^{(i)}(x) \right] \\ + \int_{x=-\infty}^{x=+\infty} \left[J[H(x)] - \lambda_j B(x) \right] d \left[S_{m_j}^{(j)}(x) - F^{(j)}(x) \right],$$

$$\begin{aligned}
 (5.17) \quad &= - \sum_{\substack{i=1 \\ i \neq j}}^c \left[\lambda_i \cdot \frac{1}{m_i} \sum_{k=1}^{m_i} \left\{ B(X_{ik}) - EB(X_i) \right\} \right] \\
 &+ \frac{1}{m_j} \sum_{k=1}^{m_j} \left\{ J[H(X_{jk})] - \lambda_j B(X_{jk}) - E[J[H(X_j)] - \lambda_j B(X_j)] \right\}
 \end{aligned}$$

where

$$B(x) = \int_{x_0}^x J'[H(y)] dF^{(j)}(y)$$

with x_0 determined somewhat arbitrarily, say by $H(x_0) = 1/2$; E represents the expectation and $X_1, \dots, X_c$ have the $F^{(1)}, \dots, F^{(c)}$ distributions respectively.

The c summations given by (5.17) involve independent samples of identically distributed random variables. Therefore, if we show that the first two moments of these random variables exist, then we can apply the central limit theorem, with the result that each sum when properly normalized will have normal distribution in the limit and hence the sum of c summations will have normal distribution in the limit.

First, to turn our attention to moments, note that by assumption 4 of lemma 5.1 and $dF^{(j)} \leq (1/\lambda_0) dH$,

$$|B(x)| \leq K \cdot [H(x)[1-H(x)]]^{-(1/2)+\delta}$$

and proceeding as in [2], for any δ' such that $(2+\delta')(-1/2+\delta) > -1$

$$E_{F^{(i)}}[|B(X)|]^{2+\delta'} \leq K; \quad i=1, \dots, j-1, j+1, \dots, c.$$

Since

$$|J(H(x)) - \lambda_j B(x)| \leq K[H(x)(1-H(x))]^{-(1/2)+\delta}$$

the existence of $2 + \delta$ absolute moments of all the terms in equation (5.17) follows.

To compute the variance of $B_{1N} + B_{2N}$, note that

$$\begin{aligned} & -\lambda_i \int_{-\infty}^{+\infty} B(x) d \left[S_{m_i}^{(i)}(x) - F^{(i)}(x) \right] \\ &= \lambda_i \int_{-\infty}^{+\infty} \left[S_{m_i}^{(i)}(x) - F^{(i)}(x) \right] J^i[H(x)] dF^{(j)}(x), \\ & \qquad i=1, \dots, j-1, \quad j+1, \dots, c, \end{aligned}$$

has mean zero and variance

$$\begin{aligned} (5.19) \quad & E \left\{ \lambda_i \int_{-\infty}^{+\infty} \left[S_{m_i}^{(i)}(x) - F^{(i)}(x) \right] J^i[H(x)] dF^{(j)}(x) \right\}^2 \\ &= E \left\{ \lambda_i^2 \int_{-\infty}^{+\infty} \int_{-\infty}^{+\infty} \left[S_{m_i}^{(i)}(x) - F^{(i)}(x) \right] \left[S_{m_i}^{(i)}(y) - F^{(i)}(y) \right] \right. \\ & \qquad \left. J^i[H(x)] J^i[H(y)] dF^{(j)}(x) dF^{(j)}(y) \right\} \\ &= \frac{2\lambda_i}{N} \iint_{-\infty < x < y < \infty} F^{(i)}(x) \left[1 - F^{(i)}(y) \right] J^i[H(x)] J^i[H(y)] \\ & \qquad dF^{(j)}(x) dF^{(j)}(y), \\ & \qquad i=1, \dots, j-1, \quad j+1, \dots, c. \end{aligned}$$

Note that the application of Fubini's theorem permits the interchange of integral and expectation.

By a similar argument, the variance of

$$\begin{aligned} & \int_{-\infty}^{+\infty} \left[J(H)(x) - \lambda_j B(x) \right] d \left[S_{m_j}^{(j)}(x) - F^{(j)}(x) \right] \\ &= - \sum_{\substack{i=1 \\ i \neq j}}^c \lambda_i \int_{-\infty}^{+\infty} \left[S_{m_j}^{(j)}(x) - F^{(j)}(x) \right] J'(H)(x) dF^{(i)}(x) \end{aligned}$$

is given by

$$\begin{aligned} (5.20) \quad & \frac{2}{N\lambda_j} \sum_{\substack{i=1 \\ i \neq j}}^c \lambda_i^2 \iint_{-\infty < x < y < \infty} F^{(j)}(x) [1 - F^{(j)}(y)] J'[H(x)] J'[H(y)] \\ & \quad dF^{(i)}(x) dF^{(i)}(y) \\ & + \frac{1}{N\lambda_j} \sum_{\substack{i,k=1 \\ i \neq k \\ i \neq j \\ k \neq j}}^c \lambda_i \lambda_k \iint_{-\infty < x < y < \infty} F^{(j)}(x) [1 - F^{(j)}(y)] J'[H(x)] J'[H(y)] \\ & \quad dF^{(i)}(x) dF^{(k)}(y) \\ & + \frac{1}{N\lambda_j} \sum_{\substack{i,k=1 \\ i \neq k \\ i \neq j \\ k \neq j}}^c \lambda_i \lambda_k \iint_{-\infty < y < x < \infty} F^{(j)}(y) [1 - F^{(j)}(x)] J'[H(x)] J'[H(y)] \\ & \quad dF^{(i)}(x) dF^{(k)}(y). \end{aligned}$$

Adding the c terms given by (5.19) and (5.20) we obtain the variance result stated in (5.7).

Thus we have shown that $B_{1N} + B_{2N}$ is the sum of c independent terms, each of which has mean zero and finite absolute $2 + \delta'$ moments. Hence sub-lemma 5.1 follows.

We shall now extend the proof of the above lemma to the case where $F^{(1)}, \dots, F^{(c)}$ and $\lambda_1, \dots, \lambda_c$ are not fixed. We want to find a set of sufficient conditions under which the asymptotic normality holds uniformly with respect to $F^{(1)}, \dots, F^{(c)}$ and $\lambda_1, \dots, \lambda_c$. For this we need the following theorem of Esseen [6], p. 43.

Theorem (Esseen) 5.1. Let $X_1, \dots, X_n$ be independent observations from a population with mean zero, variance σ^2 and finite absolute $2 + \delta'$ moments $\beta_{2+\delta'}$, $0 < \delta' \leq 1$, then

$$|F^* - \Phi^*| < c(\delta') \left| \frac{\rho_{2+\delta'}}{n^{\delta'/2}} + \frac{\rho_{2+\delta'}^{1/\delta'}}{n^{1/2}} \right|$$

where F^* is the c.d.f. of $\bar{X}$, Φ^* is the approximating normal c.d.f., $c(\delta')$ is a finite positive constant only depending on δ' and $\rho_{2+\delta'} = \beta_{2+\delta'}/\sigma^{2+\delta'}$. (If $\delta' = 1$, then $|F^* - \Phi^*| < c(\delta')\rho_3/\sqrt{n}$).

To apply this theorem in our situation, it suffices, since we have shown that the A term is finite and the C terms are uniformly $o_p(N^{-(1/2)})$, to prove the uniform convergence of $B_{1N} + B_{2N}$. For this it suffices to bound $\rho_{2+\delta'} = \beta_{2+\delta'}/\sigma^{2+\delta'}$ for $B(X_1), \dots, B(X_c)$. Since in the above lemma we already bounded the absolute $2 + \delta'$ moments, all that is required is to bound the variances of $B(X_1), \dots, B(X_c)$ away from zero. Thus we have

Corollary 5.1. If conditions 1 to 4 of lemma 5.1 are satisfied, and $F^{(1)}$ and λ_1 , $i=1, \dots, c$ (where $0 < \lambda_0 \leq \lambda_1, \dots, \lambda_c \leq 1 - \lambda_0 < 1$ holds for some fixed

$\lambda_0 \leq 1/c$) are restricted to a set for which the variances of
 $B(X_1), \dots, B(X_c)$ are bounded away from zero, then the asymptotic
normality holds uniformly with respect to $F^{(1)}, \dots, F^{(c)}$ and
 $\lambda_1, \dots, \lambda_c$.

Next we prove

Lemma 5.2. Under the assumptions of lemma 5.1, the random
vector $\sqrt{N}(T_{N,j} - \mu_{N,j}; \dots; T_{N,c} - \mu_{N,c})$ has a limiting normal
distribution.

Proof. The difference $\sqrt{N}(T_{N,j} - \mu_{N,j}) - \sqrt{N}(B_{1N}^{(j)} + B_{2N}^{(j)})$,
 where $B_{1N}^{(j)} + B_{2N}^{(j)}$ is the " $B_{1N} + B_{2N}$ " term for the j th
 component $T_{N,j} - \mu_{N,j}$, tends to zero in probability and so, by
 a well known theorem ([3], p. 299), the vectors
 $\sqrt{N}(T_{N,j} - \mu_{N,j}; \dots; T_{N,c} - \mu_{N,c})$ and $\sqrt{N}(B_{1N}^{(1)} + B_{2N}^{(1)}; \dots;$
 $B_{1N}^{(c)} + B_{2N}^{(c)})$ possess the same limiting distributions. Now since
 the j th component $B_{1N}^{(j)} + B_{2N}^{(j)}$ can be expressed as

$$\sum_{i=1}^c \left\{ \frac{1}{m_i} \sum_{\alpha=1}^{m_i} B_{ij}^*(X_{i\alpha}) \right\}, \text{ the proof of the lemma follows by apply-}$$

ing the Central Limit Theorem to each of the c independent

$$\text{vectors } \frac{1}{m_i} \sum_{\alpha=1}^{m_i} [B_{i1}^*(X_{i\alpha}), B_{i2}^*(X_{i\alpha}), \dots, B_{ic}^*(X_{i\alpha})]; i=1, \dots, c.$$

6. The Covariance of two B-Statistics. By definition

(6.1)

$$\begin{aligned} \text{Cov}(B_{1N}^{(j)} + B_{2N}^{(j)}, B_{1N}^{(j')} + B_{2N}^{(j')}) &= E(B_{1N}^{(j)} + B_{2N}^{(j)})(B_{1N}^{(j')} + B_{2N}^{(j')}) \\ &= E(B_{1N}^{(j)} B_{2N}^{(j')}) + E(B_{2N}^{(j)} B_{1N}^{(j')}) + E(B_{2N}^{(j)} B_{2N}^{(j')}) \end{aligned}$$

... ..
... ..
... ..
... ..

... ..
... ..

... ..
... ..
... ..

... ..
... ..
... ..

... ..
... ..
... ..
... ..

... ..

... ..

... ..
... ..

... ..

where

$$(6.2) \quad B_{1N}^{(j')} = \int_{[x: 0 < H(x) < 1]} J[H(x)] \left[S_{m_{j'}}^{(j')}(x) - F^{(j')}(x) \right]$$

$$(6.3) \quad B_{2N}^{(j')} = \int_{[y: 0 < H(y) < 1]} [H_N(y) - H(y)] J'[H(y)] dF^{(j')}(y)$$

and $B_{1N}^{(j)}$ and $B_{2N}^{(j)}$ are given by (5.9) and (5.10) respectively.

Now integrating $B_{1N}^{(j)}$ by parts and using the facts that

$$\int_{-\infty}^{+\infty} d \left[S_{m_j}^{(j)}(x) - F^{(j)}(x) \right] = 0$$

$$dH(x) = \sum_{i=1}^c \lambda_i dF^{(i)}(x)$$

and

$$H_N(y) - H(y) = \sum_{r=1}^c \lambda_r \left[S_{m_r}^{(r)}(y) - F^{(r)}(y) \right]$$

routine computations yield, for $j \neq j'$,

$$B_{1N}^{(j)} B_{2N}^{(j')} = - \sum_{i=1}^c \sum_{r=1}^c \lambda_i \lambda_r \iint_{x=-\infty, y=-\infty}^{x=+\infty, y=+\infty} \left[S_{m_j}^{(j)}(x) - F^{(j)}(x) \right] \left[S_{m_r}^{(r)}(y) - F^{(r)}(y) \right] J'[H(x)] J'[H(y)] dF^{(i)}(x) dF^{(j')}(y).$$

Therefore,

$$(6.4) \quad E(B_{1N}^{(j)} B_{2N}^{(j')}) = \frac{1}{N} \sum_{i=1}^c \lambda_i \iint_{-\infty < x < y < \infty} F^{(j)}(x) [1 - F^{(j)}(y)] J'[H(x)] J'[H(y)] dF^{(i)}(x) dF^{(j')}(y).$$

$$- \frac{1}{N} \sum_{i=1}^c \lambda_i \iint_{-\infty < y < x < \infty} F^{(j)}(y) [1 - F^{(j)}(x)] J'[H(x)] J'[H(y)] \\ dF^{(i)}(x) dF^{(j')}(y).$$

Proceeding analogously

$$(6.5) \quad E(B_{2N} B'_{1N}) \\ = - \frac{1}{N} \sum_{i=1}^c \lambda_i \iint_{-\infty < x < y < \infty} F^{(j')}(x) [1 - F^{(j')}(y)] \cdot J'[H(x)] \\ \cdot J'[H(y)] dF^{(i)}(x) dF^{(j)}(y) \\ - \frac{1}{N} \sum_{i=1}^c \lambda_i \iint_{-\infty < y < x < \infty} F^{(j')}(y) [1 - F^{(j')}(x)] \cdot J'[H(x)] \\ \cdot J'[H(y)] dF^{(i)}(x) dF^{(j)}(y)$$

and

$$(6.6) \quad E(B_{2N} B'_{2N}) = \frac{1}{N} \sum_{i=1}^c \lambda_i \iint_{-\infty < x < y < \infty} F^{(i)}(x) [1 - F^{(i)}(y)] \\ \cdot J'[H(x)] \cdot J'[H(y)] dF^{(j)}(x) dF^{(j')}(y) \\ + \frac{1}{N} \sum_{i=1}^c \lambda_i \iint_{-\infty < y < x < \infty} F^{(i)}(y) [1 - F^{(i)}(x)] \\ \cdot J'[H(x)] \cdot J'[H(y)] dF^{(j)}(x) dF^{(j')}(y).$$

Thus

$$(6.7) \quad N \cdot \text{Cov}(T_{N,j} - \mu_{N,j}, T_{N,j'} - \mu_{N,j'})$$

$$\begin{aligned}
&= - \sum_{i=1}^c \lambda_i \left[\iint_{-\infty < x < y < \infty} F^{(j)}(x) [1 - F^{(j)}(y)] \cdot J'[H(x)] \right. \\
&\quad \left. \cdot J'[H(y)] dF^{(i)}(x) dF^{(j')}(y) \right. \\
&+ \iint_{-\infty < y < x < \infty} F^{(j)}(y) [1 - F^{(j)}(x)] \cdot J'[H(x)] \cdot J'[H(y)] \\
&\quad \left. \cdot dF^{(i)}(x) dF^{(j')}(y) \right] \\
&- \sum_{i=1}^c \lambda_i \left[\iint_{-\infty < x < y < \infty} F^{(j')}(x) [1 - F^{(j')}(y)] \cdot J'[H(x)] \cdot J'[H(y)] \right. \\
&\quad \left. \cdot dF^{(i)}(x) dF^{(j)}(y) \right. \\
&+ \iint_{-\infty < y < x < \infty} F^{(j')}(y) [1 - F^{(j')}(x)] \cdot J'[H(x)] \cdot J'[H(y)] \\
&\quad \left. \cdot dF^{(i)}(x) dF^{(j)}(y) \right] \\
&+ \sum_{i=1}^c \lambda_i \left[\iint_{-\infty < x < y < \infty} F^{(i)}(x) [1 - F^{(i)}(y)] \cdot J'[H(x)] \cdot J'[H(y)] \right. \\
&\quad \left. \cdot dF^{(j)}(x) dF^{(j')}(y) \right. \\
&+ \iint_{-\infty < y < x < \infty} F^{(i)}(y) [1 - F^{(i)}(x)] \cdot J'[H(x)] \cdot J'[H(y)] \\
&\quad \left. \cdot dF^{(j)}(x) dF^{(j')}(y) \right], \quad j \neq j'.
\end{aligned}$$

Combining the material of the previous two sections produces

Theorem 6.1. Under the assumptions of lemma 5.1, the
random vector $T = (\sqrt{N}(T_{N,1} - \mu_{N,1}), \dots, \sqrt{N}(T_{N,c} - \mu_{N,c}))$
has a limiting normal distribution with zero mean vector
and variance - covariances given by limiting forms of
(5.7) and (6.7) respectively as $N \rightarrow \infty$.

Remark. The following theorem gives a simple sufficient condition under which conditions 1, 2, and 3 of lemma 5.1 hold.

Theorem 6.2. If $J_N(i/N)$ is the expectation of the
i-th order statistic of a sample of size N from a
population whose cumulative distribution function is the
inverse function of J

$|J^{(1)}[H(x)]| \leq K[H(1-H)]^{-1-(1/2)+\delta}$ for $i = 0, 1, 2$;
for some $\delta > 0$ and a.a. x , then

$$(1) \quad \lim_{N \rightarrow \infty} J_N(H) = J(H).$$

$$(11) \quad J_N(1) = o(\sqrt{N}).$$

$$(111) \quad \int_{I_N} [J_N(H_N) - J(H_N)] dS_{m_j}^{(j)}(x) = o(N^{-(1/2)}); \quad j=1, \dots, c.$$

Remark 1. The condition $|J^{(i)}[H(x)]| \leq K[H(1-H)]^{-i-(1/2)+\delta}$ a.a. x is weaker than the condition $|J^{(i)}(H)| \leq K[H(1-H)]^{-i-(1/2)+\delta}$ used by Chernoff and Savage [2], otherwise theorem 6.2 is the generalization of the latter's theorem 2.

Remark 2. With the use of this theorem, it is easy to verify that if $J_N(i/N)$ is the expected value of the i th order statistic of a sample of size N from (i) the standard normal distribution, (ii) the logistic distribution, (iii) the double exponential distribution, (iv) the exponential distribution, then the vector $(T_{N,1}; \dots; T_{N,c})$ has a limiting normal distribution.

7. The limiting distribution of $\mathcal{L}$ under Pitman's shift alternatives. From this section onward, we assume that $m_1, \dots, m_c$ are nondecreasing functions of a natural number n that tends to infinity. The dependence on n is indicated when necessary, by writing $m_i(n)$, $\mu_{N,i}(n)$, etc. For convenience, it is assumed that, for all i ,

$$\lim_{n \rightarrow \infty} \frac{m_i(n)}{n} = s_i$$

exists, and there exist two constants a and b such that $0 < a < s_i < b < \infty$.

In subsequent analysis, we shall concern ourselves with a sequence of admissible alternative hypothesis H_n^P which specifies that for each $i=1, \dots, c$; $F^{(i)}(x) = F(x+\theta_i/\sqrt{n})$ with $f \in \Omega$ but not specified further, and for some pair (i, j) , $\theta_i \neq \theta_j$. The letter n is used to index a sequence of situations in which H_n^P is the true hypothesis. Limiting probability distribution of $\mathcal{L}$ will then be found as $n \rightarrow \infty$.

We first prove the following

Theorem 7.1. If

(1) for all i ,

$$\lim_{n \rightarrow \infty} \frac{m_i(n)}{n} = s_i$$

exists,

(2) conditions (1) to (4) of lemma 5.1 are satisfied,

(3) $F^{(j)}(x) = F(x+\theta_j/\sqrt{n})$

so that for each index n , the hypotheses H_n^P are valid, then the random vector $[\sqrt{m_1}(T_{N,1}-\mu_{N,1}), \dots, \sqrt{m_c}(T_{N,c}-\mu_{N,c})]$ has a limiting normal distribution with zero means and covariance matrix whose (j, j') th term is

$$(7.1) \quad (\delta_{jj'} - \frac{\sqrt{s_j s_{j'}}}{\sum_{i=1}^c s_i}) A^2$$

where

$$(7.2) \quad A^2 = \int_0^1 J^2(x) dx - \left(\int_0^1 J(x) dx \right)^2$$

and the limit holds uniformly in s_1 provided

$$0 < a < s_1 < b < \infty; i=1, \dots, c.$$

Proof. From equation (5.7)

$$(7.3) \quad \lim_{n \rightarrow \infty} N \cdot \sigma_{N,j}^2 = \left[\sum_{\substack{i=1 \\ i \neq j}}^c s_i + \frac{1}{s_j} \sum_{\substack{i=1 \\ i \neq j}}^c s_i^2 + \frac{1}{2s_j} \sum_{\substack{i,k=1 \\ i \neq k \\ i \neq j \\ k \neq j}}^c s_i s_k \right] I_1 / \sum_{r=1}^c s_r$$

$$+ \frac{1}{2s_j} \left(\sum_{\substack{i,k=1 \\ i \neq k \\ i \neq j \\ k \neq j}}^c s_i s_k \right) I_2 / \sum_{r=1}^c s_r$$

where

$$(7.4) \quad I_1 = 2 \iint_{0 < x < y < 1} x(1-y) J'(x) J'(y) dx dy,$$

$$(7.5) \quad = \int_0^1 J^2(x) dx - \left(\int_0^1 J(x) dx \right)^2$$

and

$$(7.6) \quad I_2 = 2 \iint_{0 < y < x < 1} y(1-x) J'(x) J'(y) dx dy,$$

$$(7.7) \quad = \int_0^1 J^2(x) dx - \left(\int_0^1 J(x) dx \right)^2.$$

Thus, omitting the routine algebra,

$$\lim_{n \rightarrow \infty} N \cdot \sigma_{N,j}^2 = \left(-1 + \frac{\sum_{i=1}^c s_i}{s_j} \right) A^2.$$

Similarly, from equation (6.7),

$$\lim_{n \rightarrow \infty} N \text{Cov}(T_{N,j} - \mu_{N,j}, T_{N,j'} - \mu_{N,j'}) = -A^2.$$

Hence using Theorem 6.1, we obtain the desired result.

Denoting $\sqrt{m_j}(T_{N,j} - \mu_{N,j})/A$ by W_j , it now follows that the random vector $W = (W_1, \dots, W_c)$ has a limiting normal distribution with zero mean vector and with covariance matrix whose (j, j') th term is

$$\left(\delta_{jj'} - \frac{\sqrt{s_j s_{j'}}}{\sum_{i=1}^c s_i} \right).$$

We now make the analysis of variance transformation

$$S_0 = \sum_{i=1}^c \sqrt{e_i} W_i, \quad \text{where } e_i = \frac{s_i}{\sum_{i=1}^c s_i}$$

$$S_i = \sum_{i'=1}^c a_{ii'} W_{i'}, \quad ; \quad i=1, 2, \dots, c-1$$

where the a 's are chosen to make the transformation orthogonal.

It follows that $\sum_{i=1}^c W_i^2$ is asymptotically chi-square with $c-1$ degrees of freedom.

Now recalling that

$$W_i = \sqrt{m_i} \left[T_{N,i} - \mu_{N,i}(\theta) \right] / A$$

and letting

$$r_i = \sqrt{m_i} \left[\mu_{N,i}(\theta) - \mu_{N,i}(0) \right] / A$$

we write $\mathcal{L}$ as

$$\mathcal{L} = \sum_{i=1}^c (W_i + r_i)^2$$

and this has the same limiting distribution as

$$\mathcal{L}^* = \sum_{i=1}^c (W_i + r_i^*)^2$$

where $r_i^* = \lim_{n \rightarrow \infty} r_i$ reduces to

$$r_i^* = \lim_{n \rightarrow \infty} \sqrt{m_i} \left[\int_{-\infty}^{+\infty} \left[J \left\{ \sum_{\alpha=1}^c \lambda_{\alpha} F(x + \frac{\theta_{\alpha} - \theta_i}{\sqrt{n}}) \right\} - J \left\{ F(x) \right\} \right] dF(x) \right] / A.$$

We assume that the above limit exists and is finite. Noting that $\sum_{i=1}^c \sqrt{s_i} W_i = 0$ and $\sum_{i=1}^c \sqrt{s_i} r_i^* = 0$, it follows from a theorem of Mann and Wald [20] that

Theorem 7.5. Suppose that for all i , $\lim_{n \rightarrow \infty} m_i/n = s_i$ exists and is positive. Then under the hypothesis H_n^P , if for any real numbers $t_1, \dots, t_c$,

$$\lim_{n \rightarrow \infty} \sqrt{m_i} \left[\int_{-\infty}^{+\infty} \left[J \left\{ \sum_{i=1}^c \lambda_i F(x + \frac{t_i}{\sqrt{n}}) \right\} - J \left\{ F(x) \right\} \right] dF(x) \right] / A$$

exists and is finite, then for $n \rightarrow \infty$, the limiting distribution of the statistic $\mathcal{L}$ is $\chi^2_{c-1}(\lambda^L(H_n^P))$ where $\lambda^L(H_n^P)$ is the noncentrality parameter given by

$$(7.8) \quad \lambda^{\mathcal{L}}(H_n^P) = \sum_{j=1}^c \left[\lim_{n \rightarrow \infty} \sqrt{m_j} \int_{-\infty}^{+\infty} \left[J \left\{ \sum_{\alpha=1}^c \lambda_{\alpha} F(x + \frac{\theta_{\alpha} - \theta_j}{\sqrt{n}} \right\} - J \left\{ F(x) \right\} \right] dF(x) \right]^2 / A^2.$$

Remark. If the function J is such that $J(u) = u$, then from (7.8), letting $m_j = n.s_j$, we obtain for $\lambda^{\mathcal{L}}(H_n^P)$ the expression

$$\frac{12}{\left(\sum_{i=1}^c s_i\right)^2} \sum_{j=1}^c s_j \left(\sum_{\alpha=1}^c s_{\alpha} \lim_{n \rightarrow \infty} \int_{x=-\infty}^{x=+\infty} \sqrt{n} \left\{ F(x + \frac{\theta_{\alpha} - \theta_j}{\sqrt{n}}) - F(x) \right\} dF(x) \right)^2$$

which is the noncentrality parameter $\lambda^H(H_n^P)$ of the Kruskal-Wallis H test. (See Andrews [1], p. 726).

In many situations, the noncentrality parameter $\lambda^{\mathcal{L}}$ can be computed easily with the aid of the following lemma which, though stated in a form appropriate to our purpose, is due to Hodges and Lehmann [11].

Lemma 7.2 (Hodges-Lehmann). If

(1) F is a continuous cumulative distribution function, differentiable in each of the open intervals $(-\infty, a_1)$, (a_1, a_2) , $\dots, (a_{s-1}, a_s)$, (a_s, ∞) and the derivative of F is bounded in each of these intervals and

(ii) the function $\frac{d}{dx} J[F(x)]$ is bounded as $x \rightarrow \pm \infty$

then

$$(7.9) \quad \lim_{n \rightarrow \infty} \sqrt{n} \int_{-\infty}^{+\infty} \left[J \left\{ \sum_{\alpha=1}^c \lambda_{\alpha} F(x + \frac{\theta_{\alpha} - \bar{\theta}}{\sqrt{n}}) \right\} - J \{ F(x) \} \right] dF(x) \\ = \frac{1}{\sum_{\alpha=1}^c s_{\alpha}} \sum_{\alpha=1}^c s_{\alpha} (\theta_{\alpha} - \bar{\theta}) \int_{-\infty}^{+\infty} \frac{d}{dx} J \{ F(x) \} dF(x).$$

The proof of this lemma follows by the methods used in section 3 and 4 of Hodges-Lehmann (1961).

In case the conditions of lemma 7.2 are satisfied, then

$$(7.10) \quad \lambda^L(H_n^P) = \sum_{\alpha=1}^c s_{\alpha} (\theta_{\alpha} - \bar{\theta})^2 \left(\int_{-\infty}^{+\infty} \frac{d}{dx} J[F(x)] f(x) dx \right)^2 / A^2$$

where

$$(7.11) \quad \bar{\theta} = \sum_{\alpha=1}^c s_{\alpha} \theta_{\alpha} / \sum_{\alpha=1}^c s_{\alpha}$$

and A^2 is defined in (7.2).

8. Asymptotic relative efficiency. The concept of asymptotic relative efficiency of one test with respect to another is due to Pitman. An exposition of his work, together with some extensions is presented by Noether [27].

Theorem 8.1. If $m_1 = n \cdot s_1$, and if the distribution function F is such that

$$(1) \quad \lim_{n \rightarrow \infty} \sqrt{n} \int_{-\infty}^{+\infty} \left[F(x + \frac{\theta}{\sqrt{n}}) - F(x) \right] dF(x) \text{ exists}$$

$$(2) \quad \lim_{n \rightarrow \infty} \sqrt{n} \int_{-\infty}^{\infty} \left[J \left\{ \frac{1}{\sum_{i=1}^c s_i} \sum_{\alpha=1}^c s_{\alpha} F \left(x + \frac{\theta_{\alpha} - \theta_j}{\sqrt{n}} \right) \right\} - J[F(x)] \right] dF(x) / A$$

exists

then the asymptotic relative efficiency of the H test with respect to an arbitrary L test for testing the hypothesis H_0 against H_n^P is given by

$$(8.1) \quad e_{H,L}^P(F(x))$$

$$= \frac{12 \sum_{\alpha=1}^c s_{\alpha} \left(\sum_{i=1}^c s_i \lim_{n \rightarrow \infty} \int_{-\infty}^{+\infty} \sqrt{n} \left[F \left(x + \frac{\theta_i - \theta_{\alpha}}{\sqrt{n}} \right) - F(x) \right] dF(x) \right)^2 A^2}{\left(\sum_{j=1}^c s_j \right)^2 \sum_{\alpha=1}^c s_{\alpha} \left(\lim_{n \rightarrow \infty} \int_{-\infty}^{+\infty} \sqrt{n} \left[J \left\{ \frac{1}{\sum_{j=1}^c s_j} \sum_{i=1}^c s_i F \left(x + \frac{\theta_i - \theta_{\alpha}}{\sqrt{n}} \right) \right\} - J[F(x)] \right] dF(x) \right)^2}$$

The proofs of the above theorem follows by taking the ratio of the two noncentrality factors after the alternatives have been equated.

Corollary 8.1. If in addition to the hypotheses of theorem 8.1, the hypotheses of lemma 7.2 are satisfied, then

$$(8.2) \quad e_{H,L}^P(F(x)) = 12A^2 \left(\frac{\int_{-\infty}^{+\infty} f^2(x) dx}{\int_{-\infty}^{+\infty} \frac{d}{dx} J[F(x)] f(x) dx} \right)^2$$

where f is the density of F.

Here $e_{H,L}^P$ does not depend upon c, α, β , and is a function of F only.

It may be remarked that (8.2) agrees with the results found by Chernoff-Savage [2] and Hodges-Lehmann [11] for the two-sample case, and hence the results of this paper as well as those of [2] apply directly to the c-sample problem.

The asymptotic relative efficiency of the classical $\mathcal{F}$ test with respect to an arbitrary $\mathcal{L}$ test is contained in the following

Theorem 8.2. If

(i) for all i , $\lim_{n \rightarrow \infty} \frac{m_i(n)}{n} = s_i$ exists and is positive,

(ii) the distribution function $F(x)$ satisfies the assumptions of lemma 7.2, and

(iii) $\int_{-\infty}^{+\infty} x^2 dF(x) - \left(\int_{-\infty}^{+\infty} x dF(x) \right)^2 = \sigma^2$ exists,

then, the asymptotic relative efficiency of the classical $\mathcal{F}$
test with respect to an arbitrary $\mathcal{L}$ test for testing the
hypothesis H_0 against H_n^P is

$$(8.3) \quad e_{\mathcal{F}, \mathcal{L}}^P(F(x)) = \frac{A^2}{\sigma^2} \left(\frac{1}{\int_{-\infty}^{+\infty} \frac{d}{dx} J[F(x)] dF(x)} \right)^2$$

Proof. The $\mathcal{F}$ statistic is defined as

$$\mathcal{F} = \frac{\frac{1}{c-1} \sum_{i=1}^c m_i (X_{i.} - \bar{X})^2}{\frac{1}{N-c} \sum_{i=1}^c \sum_{j=1}^{m_i} (X_{ij} - X_{i.})^2}$$

where $X_{i.} = \sum_{j=1}^{m_i} X_{ij}/m_i$ and $\bar{X} = \sum_{i=1}^c \sum_{j=1}^{m_i} X_{ij}/N$. It has been

shown by Andrews [1] that under the hypothesis H_n^P , this has a limiting noncentral chi-square distribution with $c-1$ degrees of freedom and noncentrality parameter $\lambda^{\frac{1}{2}}(H_n^P)$ given by

$$(8.5) \quad \lambda^{\frac{1}{2}}(H_n^P) = \sum_{i=1}^c s_i [(\theta_i - \bar{\theta})/\sigma]^2.$$

Now proceeding by standard arguments, the proof follows.

In particular, when $J = \Phi^{-1}$, where Φ is the standard cumulative normal distribution function having the density ϕ ,

$$(8.6) \quad e_{\mathcal{L}, \frac{1}{2}}^P(F(x)) = \sigma^2 \left(\int_{-\infty}^{+\infty} \frac{f^2(x) dx}{\phi \{ \Phi^{-1}[F(x)] \}^2} \right)^2,$$

which is known to be the asymptotic efficiency of the two sample normal scores test with respect to the student's t-test and is always ≥ 1 . When $F(x)$ is normal distribution function, this is 1. See in this connection Chernoff-Savage [2] and Hodges-Lehmann [11].

2. Acknowledgment. I wish to express my sincere thanks to Professor Erich L. Lehmann for proposing this investigation and for the generous help and guidance during its progress.

APPENDIX

10. Higher order terms. Before we prove that the c terms of lemma 5.1 are uniformly of higher order, we state the following elementary results which are used repeatedly. (Also in this connection see Chernoff and Savage [2].)

10A. Elementary results.

$$1. \quad H \geq \lambda_1 F^{(1)} \geq \lambda_0 F^{(1)}; \quad i=1, \dots, c.$$

$$2. \quad 1 - F^{(1)} \leq \frac{1-H}{\lambda_1} \leq \frac{1-H}{\lambda_0}; \quad i=1, \dots, c.$$

$$3. \quad F^{(1)}(1-F^{(1)}) \leq \frac{H(1-H)}{\lambda_1^2} \leq \frac{H(1-H)}{\lambda_0^2}; \quad i=1, \dots, c.$$

$$4. \quad dH \geq \lambda_1 dF^{(1)} \geq \lambda_0 dF^{(1)}; \quad i=1, \dots, c.$$

5. Let (a_N, b_N) be the interval S_{N_ϵ} where

$$(10.1) \quad S_{N_\epsilon} = [x: H(1-H) > \eta_\epsilon \lambda_0 / N],$$

when η_ϵ can be chosen independent of $F^{(1)}$ and λ_1 ; $i=1, \dots, c$, such that

$$(10.2) \quad P[X_{1j} \in S_{N_\epsilon}; i=1, \dots, c; j=1, \dots, m_1] \geq 1 - \epsilon.$$

10B. Detailed consideration of the c -terms of lemma 5.1.

First, let us consider

(10.3)

$$c_{1N} = \lambda_1 \int_{0 < H < 1} \left[S_{m_1}^{(i)}(x) - F^{(i)}(x) \right] J'[H(x)] d \left[S_{m_j}^{(j)}(x) - F^{(j)}(x) \right];$$

$$i=1, \dots, j-1, j+1, \dots, c$$

$$= \lambda_1 \left[c_{1N}^{(i)} + c_{2N}^{(i)} \right]; \quad i=1, \dots, c; \quad i \neq j,$$

where

(10.4)

$$c_{1N}^{(i)} = \int_{S_{N_\epsilon}} \left[S_{m_1}^{(i)}(x) - F^{(i)}(x) \right] J'[H(x)] d \left[S_{m_j}^{(j)}(x) - F^{(j)}(x) \right]$$

$$i=1, \dots, c; \quad i \neq j,$$

and

(10.5)

$$c_{2N}^{(i)} = \int_{S_{N_\epsilon}} \left[S_{m_1}^{(i)}(x) - F^{(i)}(x) \right] J'[H(x)] d \left[S_{m_j}^{(j)}(x) - F^{(j)}(x) \right];$$

$$i=1, \dots, c; \quad i \neq j.$$

First note that

(10.6)

$$E(c_{1N}^{(i)}) = E[E(c_{1N}^{(i)} | X_{j1}, \dots, X_{jm_j})] = 0; \quad i=1, \dots, c; \quad i \neq j.$$

 

7.

8.

 

Next

$$\begin{aligned} [C_{1N}^{(i)}]^2 &= 2 \iint_{\substack{x, y \in S_{N_\epsilon} \\ x < y}} \left[S_{m_i}^{(i)}(x) - F^{(i)}(x) \right] \left[S_{m_i}^{(i)}(y) - F^{(i)}(y) \right] J'[H(x)] J'[H(y)] \\ &\quad d \left[S_{m_j}^{(j)}(x) - F^{(j)}(x) \right] d \left[S_{m_j}^{(j)}(y) - F^{(j)}(y) \right] \\ &\quad + \frac{1}{m_j} \int_{x \in S_{N_\epsilon}} \left[S_{m_i}^{(i)}(x) - F^{(i)}(x) \right]^2 \left[J'[H(x)] \right]^2 dS_{m_j}^{(j)}(x); \end{aligned}$$

$$i=1, \dots, c; \quad i \neq j.$$

Therefore,

$$\begin{aligned} (10.7) \quad E(C_{1N}^{(i)})^2 &= E \left[E[(C_{1N}^{(i)})^2 | X_{j1}, \dots, X_{jm_j}] \right] \\ &\quad - \frac{2}{m_i m_j} \iint_{\substack{x, y \in S_{N_\epsilon} \\ x < y}} F^{(i)}(x) [1 - F^{(i)}(y)] J'[H(x)] J'[H(y)] \\ &\quad \quad \quad dF^{(j)}(x) dF^{(j)}(y) \\ &\quad + \frac{1}{m_i m_j} \int_{x \in S_{N_\epsilon}} F^{(i)}(x) [1 - F^{(i)}(x)] [J'[H(x)]]^2 dF^{(j)}(x); \end{aligned}$$

$$i=1, \dots, c; \quad i \neq j$$

$$\begin{aligned} &\leq \frac{K}{N^2} \iint_{\substack{x, y \in S_{N_\epsilon} \\ x < y}} H(x) [1 - H(y)] [H(x)(1 - H(x))]^{-3/2+\delta} [H(y)(1 - H(y))]^{-3/2+\delta} \\ &\quad + \frac{K}{N^2} \int_{x \in S_{N_\epsilon}} H(x) [1 - H(x)] [H(x)(1 - H(x))]^{-3+2\delta} dH(x) dH(y) \\ &\leq \frac{K}{N^2} + \frac{K \eta_\epsilon^{-1+2\delta}}{N^{1+2\delta}} = o\left(\frac{1}{N}\right); \quad [K \text{ is generic}]. \end{aligned}$$

Hence from (10.6) and (10.7), we obtain, using the Markoff inequality,

$$(10.8) \quad \left| c_{1N}^{(1)} \right| = o_p(N^{-(1/2)}).$$

We now consider $c_{2N}^{(1)}$. Let $H_1 = H(a_N)$, $H_2 = H(b_N)$. Then from (10.1) $H_1 = 1 - H_2 < K/N$. With probability greater than $1 - \varepsilon$, there are no observations in $\bar{S}_{N\varepsilon}$ and

$$(10.9) \quad \left| c_{2N}^{(i)} \right| \leq \int_0^{H_1} F^{(i)}(x) [J'[H(x)]] dF^{(j)}(x) \\ + \int_{H_2}^1 (1-F^{(i)}(x)) [J'[H(x)]] dF^{(j)}(x)$$

$$i=1, \dots, c; i \neq j$$

$$\leq K \int_0^{H_1} \frac{H dH}{[H(1-H)]^{(3/2)-\delta}} + \int_{H_2}^1 \frac{(1-H) dH}{[H(1-H)]^{(3/2)-\delta}}$$

$$\leq K \int_0^{H_1} H^{-(1/2)+\delta} dH \leq K \frac{1}{N^{(1/2)+\delta}}.$$

Hence

$$(10.10) \quad \left| c_{2N}^{(i)} \right| = o_p(N^{-(1/2)}); \quad i=1, \dots, c; i \neq j.$$

Consequently,

$$(10.11) \quad c_{1N} = \lambda_1 \left[c_{1N}^{(1)} + c_{2N}^{(1)} \right] = o_p(N^{-(1/2)}); i=1, \dots, c; i \neq j.$$

The proof of $c_{jN} = o_p(N^{-(1/2)})$ follows first showing that

$$c_{jN} = -\frac{\lambda_j}{2} \left[c_{11N} + c_{12N} - c_{13N} \right]$$

where

$$(a) \quad C_{11N} = \int_{S_{N_\varepsilon}} \left[S_{m_j}^{(j)}(x) - F^{(j)}(x) \right]^2 J''[H(x)] dH(x),$$

$$(b) \quad C_{12N} = \int_{S_{N_\varepsilon}} \left[S_{m_j}^{(j)}(x) - F^{(j)}(x) \right]^2 J''[H(x)] dH(x),$$

$$(c) \quad C_{13N} = \frac{1}{m_j} \int J'[H(x)] dS_{m_j}^{(j)}(x)$$

and then showing that each C_{1kN} is $o_p(H^{-(1/2)})$; $k=1,2,3$.

The proofs of the above statement are omitted since they are essentially contained in the work of Chernoff and Savage [2].

Next consider

$$C_{c+1,N} = \int_{I_N} [H_N(x) - H(x)]^2 J''[\theta H_N(x) + (1-\theta)H(x)] dS_{m_j}^{(j)}(x),$$

$$0 < \theta < 1.$$

With probability $> 1 - \varepsilon$, the interval I_N can be replaced by S_{N_ε} without changing $C_{c+1,N}$. Furthermore since

$$\sup_{H_N > 0} \left| \frac{H(x)}{H_N(x)} \right| = o_p(1)$$

and

$$\sup_{H_N < 1} \left| \frac{1-H(x)}{1-H_N(x)} \right| = o_p(1),$$

for each $\varepsilon > 0$, there exists an $\eta_\varepsilon^* > 0$ such that with greater

than $1 - \epsilon$, we have for $\{x: 0 < H_N(x) < 1\}$,

$$[\theta H_N(x) + (1-\theta)H(x)][1 - \{\theta H_N(x) + (1-\theta)H(x)\}] > \eta_\epsilon^* H(x)[1-H(x)].$$

Then

$$|C_{c+1,N}| \leq (\eta_\epsilon^*)^{-(5/2)+\delta} C_{cN}$$

where

$$C_{cN} = \int_{S_{N_\epsilon}} [H_N(x) - H(x)]^2 \{H(x)[1-H(x)]\}^{-(5/2)+\delta} dS_{m_j}^{(j)}(x)$$

and

$$\begin{aligned} E(C_{cN}) &= E[E(C_{cN} | X_{j1}, \dots, X_{jm_j})] \\ &= \frac{1}{N} \int_{S_{N_\epsilon}} \sum_{i=1}^c \lambda_i F^{(i)}(1-F^{(i)}) [H(1-H)]^{-(5/2)+\delta} dF^{(j)}(x) \\ &\quad + \frac{1}{N^2} \int_{S_{N_\epsilon}} (1-F^{(j)})(1-2F^{(j)}) [H(1-H)]^{-(5/2)+\delta} dF^{(j)}(x) \\ &\leq \frac{K}{N} \int_{S_{N_\epsilon}} [H(1-H)]^{-(3/2)+\delta} dH + \frac{K}{N^2} \int_{S_{N_\epsilon}} [H(1-H)]^{-(5/2)+\delta} dH \\ &\leq \frac{K}{N(1/2)+\delta}. \end{aligned}$$

Consequently $C_{c+1,N} = o_p(N^{-(1/2)})$.

The negligibility of $C_{c+2,N}$ and $C_{c+3,N}$ follows from assumptions 2 and 3 of lemma 5.1 and the proof of the negligibility of $C_{c+4,N}$ proceeds in the same manner as given by Chernoff and Savage for the term C_{4N} and therefore is not given here.

REFERENCES

1. ANDREWS, F. C. (1954). Asymptotic behavior of some rank tests for analysis of variance. Ann. Math. Statist. 25 724-735.
2. CHERNOFF, H. and SAVAGE, I. R. (1958). Asymptotic normality and efficiency of certain non-parametric test statistics. Ann. Math. Statist. 29 972-994.
3. CRAMÉR, Harald (1946). Mathematical Methods of Statistics. Princeton University Press.
4. CRAMÉR, Harald (1937). Random Variables and Probability Distributions. Cambridge University Press.
5. DANTZIG, D. Van (1951). On the consistency and the power of Wilcoxon's two-sample test. Nederl. Akad. Wetensch. Proc. 54
6. ESSEEN, C. G. (1945). Fourier analysis of distribution functions. Acta Math. 77 1-125.
7. FRASER, D. A. S. (1956). Non-parametric Methods of Statistics. John Wiley and Sons, Inc.
8. FREUND, J. E. and ANSARI, A. R. (1957). Two-way rank sum for variances. Virginia Polytechnic Institute.
9. HANNAN, E. J. (1949). The asymptotic powers of certain tests based on multiple correlations. J. Roy. Statist. Soc. B. 11 21-219.
10. HODGES, J. L. JR. and LEHMANN, E. L. (1956). The efficiency of some non-parametric competitors of the t-test. Ann. Math. Statist. 27 324-335.

11. HODGES, J. L. JR. and LEHMANN, E. L. (1961). Comparison of the normal scores and Wilcoxon tests. Proceedings of the Fourth Berkeley Symposium on Mathematical Statistics and Probability. 1 307-317.
12. KIEFER, J. (1959). K-sample analogues of the Kolmogorov Smirnov and Cramer v. Mises tests. Ann. Math. Statist. 30 420-447.
13. KRUSKAL, W. H. (1952). A non-parametric test for the several sample problems. Ann. Math. Statist. 23 525-540.
14. KRUSKAL, W. H. and WALLIS, W. A. (1952). Use of ranks in one criterion analysis of variance. J. Amer. Statist. Assoc. 47 583-621.
15. LEHMANN, E. L. (1951). Consistency and unbiasedness of certain non-parametric tests. Ann. Math. Statist. 22 165-179.
16. LEHMANN, E. L. (1953). The power of rank tests. Ann. Math. Statist. 24 23-43.
17. LEHMANN, E. L. (1959). Testing Statistical Hypotheses. John Wiley and Sons, Inc.
18. LOEVE, M. (1955). Probability Theory. D. Van Nostrand Co., Inc.
19. MANN, H. B. and WHITNEY, D. R. (1947). On a test of whether one of two random variables is stochastically larger than the other. Ann. Math. Statist. 18 50-60.
20. MANN, H. B. and WALD, A. (1943). On stochastic limit and order relationship. Ann. Math. Statist. 14 217-226.

21. MIHUKSKI, P. W. (1961). Some problems in the asymptotic theory of testing statistical hypotheses. Efficiency of non-parametric procedures. Ph.D. Thesis, University of California.
22. MOOD, A. M. (1950). Introduction to Theory of Statistics. McGraw-Hill Publishing Co.
23. MOOD, A. M. (1954). On the asymptotic efficiency of certain non-parametric two-sample tests. Ann. Math. Statist. 25 514-522.
24. RAO, C. R. (1952). Advanced Statistical Methods in Biometric Research. John Wiley and Sons, Inc.
25. SCHEFFÉ, H. (1943). Statistical inference in the non-parametric case. Ann. Math. Statist. 14 305-332.
26. TERPESTRA, T. J. (1952). A non-parametric k-sample test and its connection with the H-test. Report S(92) (VP2) of the Statistical Department of the Mathematical Centre, Amsterdam.
27. TERRY, M. E. (1952). Some rank order tests which are most powerful against specific alternatives. Ann. Math. Statist. 23 346-366.
28. WALD, A. and WOLFOWITZ, J. (1944). Statistical tests based on permutations of the observations. Ann. Math. Statist. 15 358-372.
29. WILCOXON, F. (1945). Individual comparison by ranking methods. Biometrics Bull. 1 80-83.

30. VAN DER WAERDEN, D. L. (1953). Ein neuer Test für das
problem der Zwei Stich problem. Math. Annalen. 126
93-107.
31. VAN DER WAERDEN, B. L. (1956). The computation of the χ^2 -
distribution. Proceedings of the Third Berkeley
Symposium on Mathematical Statistics and Probability.
1 207-208.
32. NOETHER, G. E. (1954). On a theorem of Pitman. Ann. Math.
Statist. 25 514-522.
33. ANSARI, A. R. and BRADLEY, R. A. (1960). Rank sum tests
for dispersions. Ann. Math. Statist. 31 1174-1184.
34. SAVAGE, I. R. (1952). Contributions to the theory of rank
order statistics: Two-sample case. Ann. Math. Statist.
23 346-366.
35. HOEFFDING, W. (1950). Optimum non-parametric tests.
Proceedings of the Second Berkeley Symposium on
Mathematical Statistics and Probability. 83-92.

Approved Distribution List for Unclassified Technical Reports

Head, Statistics Branch Office of Naval Research Washington 25, D.C. (2)	Professor Norman Chernoff Applied Math. and Stat. Lab. Stanford University Stanford, California (1)
Commanding Officer Office of Naval Research Branch Office, Navy 100 Fleet Post Office New York, New York (2)	Professor W.G. Cochran Department of Statistics Harvard University Cambridge, Massachusetts (1)
ASTIA Document Service Center Arlington Hall Station Arlington 12, Virginia (10)	Professor Benjamin Epstein Applied Math. and Stat. Lab. Stanford University Stanford, California (1)
Technical Information Officer Naval Research Laboratory Washington 25, D.C. (6)	Professor W. Hirsch Inst. of Math. Sciences New York University New York 3, New York (1)
Institute for Defense Analyses Communications Research Division von Neumann Hall Princeton, New Jersey (1)	Dr. Paul G. Hoel Department of Mathematics University of California Los Angeles 24, California (1)
Bureau of Supplies and Accounts Code OW Department of the Navy Washington 25, D.C. (1)	Professor Harold Hotelling Associate Director Institute of Statistics University of North Carolina Chapel Hill, North Carolina (1)
Professor T.W. Anderson Department of Mathematical Statistics Columbia University New York 27, New York (1)	Professor L. Hurwicz School of Business Administration University of Minnesota Minneapolis, Minnesota (1)
Professor A.J. Birnbaum Laboratory of Statistical Research Department of Mathematics University of Washington Seattle 5, Washington (1)	Professor Leo Katz Department of Statistics Michigan State University East Lansing, Michigan (1)
Professor A.H. Bowker Applied Mathematics and Statistical Lab. Stanford University Stanford, California (1)	Professor Oscar Kempthorne Statistics Laboratory Iowa State College Ames, Iowa (1)
Professor Ralph A. Bradley Department of Stat. and Statistical Lab. Virginia Polytechnic Institute Blacksburg, Virginia (1)	Dr. Carl F Kossack I.B.M. Corp. Data Processing 2601 Main St. Houston 2, Texas (1)

Approved Distribution List (Con't.)

Professor Gerald J. Lieberman
Applied Mathematics and
Statistics Laboratory
Stanford University
Stanford, California (1)

Professor William G. Madow
Stanford Research Institute
Menlo Park, California (1)

Professor J. Neyman
Department of Statistics
University of California
Berkeley 4, California (1)

Professor Herbert Robbins
Mathematical Statistics Department
Columbia University
New York 27, New York (1)

Professor Murray Rosenblatt
Department of Mathematics
Brown University
Providence, R.I. (1)

Professor L. J. Savage
Department of Mathematics
University of Michigan
Ann Arbor, Michigan (1)

Dr. Herbert Solomon
Department of Statistics
Stanford University
Stanford, California (1)

Professor Frank Spitzer
Department of Mathematics
Cornell University
Ithaca, New York (1)

Professor S.S. Wilks
Department of Mathematics
Princeton University
Princeton, New Jersey (1)

Professor Evan J. Williams
Institute of Statistics
State College Section
North Carolina State College
Raleigh, North Carolina (1)

Professor J. Wolfowitz
Department of Mathematics
Cornell University
Ithaca, New York (1)

Professor W.H. Kruskal
Department of Statistics
The University of Chicago
Chicago 37, Illinois (1)

DATE DUE

5-10-88			
5-1-20			
GAYLORD			PRINTED IN U.S.A.

NYU
IMM-
314

c.1

Puri

NYU
IMM-
314

c.1

NYU
IMM-
314

c.1

314 Puri

Asymptotic efficiency of a
class of c-sample tests.

**N. Y. U. Courant Institute of
Mathematical Sciences**

4 Washington Place
New York 3, N. Y.